



EDINBOROUGH STREET RADIOLOGY INC. & PULMONARY FUNCTION LAB

PULMONARY FUNCTION TEST & RESPIRATORY CONSULTATION REFERRAL FORM

PATIENT INFORMATION	PHYSICIAN INFORMATION
Name: _____ DOB _____ <input type="checkbox"/> M <input type="checkbox"/> F Address: _____ City: _____ Prov: _____ Postal Code: _____ Phone (Home) _____ Cell) _____ Health Card # _____ VC _____	Name: _____ Address: _____ City: _____ Prov: _____ Postal Code: _____ Phone: _____ Fax: _____ OHIP Provider # _____ Copy To: _____ Signature: _____

REQUEST FOR:

- Consultation/Pulmonary Function Testing
- Pulmonary Function Testing only
- Consultation only

CLINICAL INFORMATION:

1. Reason for Referral:

2. Respiratory Medications: No Yes (Please List)
Current Respiratory Medications:

3. Smoking History:

4. Tests Requested (Please check)

- Full Pulmonary Function Testing** (Spirometry, flow-volume loops with bronchodilator if indicated, Lung volumes and plethysmography for airway resistance/lung diffusion capacity).
- Spirometry/Flow Volume Loops:** (including post-bronchodilator study if indicated)
- MIPS/MEPS:** (maximum inspiratory/expiratory)

APPOINTMENT DATE and TIME: _____

Please see reverse for instructions

**SMOKING SHOULD BE DISCONTINUED ON THE DAY OF TESTING.
 CAFFEINATED BEVERAGES SHOULD BE AVOIDED ON THE DAY OF TESTING.**

MEDICATIONS WHICH INTERFERE WITH PULMONARY FUNCTION TESTING

Some medications will interfere with pulmonary function testing and should not be taken prior to your test, if possible. However, if you do not think that you can go without, or you develop significant respiratory symptoms, then continue your usual routine and inform the therapist that you have taken your medications before starting your test.

MEDICATION		ABSTINENCE
Short acting bronchodilators	Salbutamol (airomir; entolin) Terbutaline (bricanyl)	8 hours
Long acting bronchodilators	Indacaterol (onbrez) Formoterol (oxeze) Salmeterol (serevent) Vilanterol	48 hours
Anticholinergics Short Acting	Ipratropium (atrovent)	24 hours
Anticholinergics Long Acting	Aclidinium (tudorza) Glycopyrronium (seebri) Tiotropium; (spiriva) Umeclidinium (incruise)	48 hours
Leukotriene receptor antagonists	Montelukast (singulair)	24 hours
Theophylline	Twice daily preparation Once daily preparation	24 hours 48 hours

Anti-inflammatory medications containing steroids do not need to be stopped. These medications include:

- Beclomethasone (QVAR)
- Budesonide (Pulmicort)
- Ciclesonide (Alvesco)
- Fluticasone (Flovent)
- Mometasone (Asmanex)
- Prednisone

However, anti-inflammatory medications which are combination medications need to be stopped according to the table above. These medications include:

- Fluticasone-vilanterol (Breo)
- Fluticasone-umeclidinium-vilanterol (Trelegy)
- Formoterol-budesonide (Symbicort)
- Formoterol-mometasone (Zenhale)
- Salmeterol-fluticasone (Advair)

NOTE: There is a \$125 charge for missed/cancelled consultation appointment without 48 business hours prior notification and a \$75 charge for missed/cancelled pulmonary function testing appointment without 48 business hours prior notification

The patient will be contacted directly to arrange an appointment. Your office will be notified the appointment date & time.